REFERENCES:

1. Aragao GF , Weintraub ST, Yoshihama T, Maruyama Y *et al*., (2006), A possible mechanism for anxiolytic and antidepressant effects of alpha- and beta-amyrin from Protium heptaphyllum (Aubl.) March, Pharmacol Biochem Behav, 85(4), 827-34.
2. Roychoudhury M, Kulkarni SK (1997), Antianxiety profile of ondansetron, a selective 5-HT3 antagonist, in a novel animal model, Methods Find Exp Clin Pharmacol, 19(2), 107-11.
3. Kasper, Braunwald, Fauci, Hauser, Longo, Jameson. Harrison's principles of internal medicine. 16th edition. McGraw-Hill Publication, Newyork, Vol-2: 2547-2552.
4. www.cnsforum.com/ imagebank/section/Brain_struc_anxiety/default.aspx
5. Curtis GC, Magee WJ, Eaton WW. Specific fears and phobias. Epidemiology and classification. Br J Psychiatry, 1998; 173:212-17.
6. Kuribara H, Weintraub ST, Yoshihama T, Maruyama Y (2003), An anxiolytic-like effect of Ginkgo biloba extract and its constituent, ginkgolide-A, in mice, J Nat Prod, 66(10), 1333-7.
7. Nogueira E, Rosa GJ, Vassilieff VS (1998), Involvement of GABA(A)-benzodiazepine receptor in the anxiolytic effect induced by hexanic fraction of Rubus brasiliensis, J Ethnopharmacol, 61(2), 119-26.
8. Peng WH, Hsieh MT, Lee YS, Lin YC, Liao J (2000), Anxiolytic effect of seed of Ziziphus jujuba in mouse models of anxiety, J Ethnopharmacol, 72(3), 435-41.
9. Schmidt-Mutter C, Pain L, Sandner G, Gobaille S, Maitre M (1998), The anxiolytic effect of gamma-hydroxybutyrate in the elevated plus maze is reversed by the benzodiazepine receptor antagonist, flumazenil, Eur J Pharmacol, 342(1), 21-7.
10. Garrett KM, Basmadjian G, Khan IA, Schaneberg BT, Seale TW (2003), Extracts of kava (Piper methysticum) induce acute anxiolytic-like behavioural changes in mice, Psychopharmacology (Berl), 170(1) 33-41.
11. Nogueira E, Rosa GJ, Vassilieff VS (1998), Involvement of GABA(A)-benzodiazepine receptor in the anxiolytic effect induced by hexanic fraction of Rubus brasiliensis, J Ethnopharmacol, 61(2), 119-26.
12. Wieronska JM, Papp M, Pilc A (2001), Effects of anxiolytic drugs on some behavioural consequences in olfactory bulbectomized rats, Pol J Pharmacol, 53(5), 517-25.

13. Przegaliński E, Tatarczyńska E, Chojnacka-Wójcik E (200), The influence of the benzodiazepine receptor antagonist flumazenil on the anxiolytic-like effects of CGP 37849 and ACPC in rats, Neuropharmacology, 39(10), 1858-64.
14. Hattesohl M, Feistel B, Sievers H, Lehnfeld R, Hegger M, Winterhoff H (2008), Extracts of Valeriana officinalis L. s.l. show anxiolytic and antidepressant effects but neither sedative nor myorelaxant properties, Phytomedicine, (1-2), 2-15.
15. Grundmann O, Nakajima J, Seo S, Butterweck V (2007), Anti-anxiety effects of Apocynum venetum L. in the elevated plus maze test, J Ethnopharmacol, 110(3), 406-11.
16. Jung JW, Ahn NY, Oh HR, Lee BK, Lee KJ, Kim SY, Cheong JH, Ryu JH (2006), Anxiolytic effects of the aqueous extract of Uncaria rhynchophylla, J Ethnopharmacol, 108(2), 193-7.
17. Boschloo L, Vogelzangs N, Van den Brink W, Smit JH, Veltman DJ, Beekman AT, Penninx BW (2012), Alcohol use disorders and the course of depressive and anxiety disorders, Br J Psychiatry.
18. Fakeye TO, Pal A, Khanuja SP (2008), Anxiolytic and sedative effects of extracts of Hibiscus sabdariffa Linn (family Malvaceae), Afr J Med Sci, 37(1), 49-54.
19. Cardoso Vilela F, Soncini R, Giusti-Paiva A (2009), Anxiolytic-like effect of Sonchus oleraceus L. in mice, J Ethnopharmacol. 124(2), 325-7.
20. Deng J, Zhou Y, Bai M, Li H, Li L (2010), Anxiolytic and sedative activities of Passiflora edulis f. Flavicarpa, J Ethnopharmacol, 128(1), 148-53.
21. Herrera-Ruiz M *et al*., (2006), Antidepressant and anxiolytic effects of hydroalcoholic extract from Salvia elegans, J Ethnopharmacol, 107(1), 53-8.
22. Kuribara H, Weintraub ST, Yoshihama T, Maruyama Y (2003), An anxiolytic-like effect of Ginkgo biloba extract and its constituent, ginkgolide-A, in mice, J Nat Prod, 66(10), 1333-7.
23. Aragao GF *et al*., (2006), A possible mechanism for anxiolytic and antidepressant effects of alpha- and beta-amyrin from Protium heptaphyllum (Aubl.) March, Pharmacol Biochem Behav, 85(4), 827-34.
24. Rabbani M, Sajjadi SE, Zarei HR (2003), Anxiolytic effects of Stachys lavandulifolia Vahl on the elevated plus-maze model of anxiety in mice, J Ethnopharmacol, 89(2-3), 271-6.

25. Hellion-Ibarrola MC *et al*., (2006), the anxiolytic-like effects of Aloysia polystachya (Griseb.) Moldenke (Verbenaceae) in mice, J Ethnopharmacol, 105(3), 400-8.
26. Tolardo R *et al*., (2010), Evaluation of behavioural and pharmacological effects of Hedyosmum brasiliense and isolated sesquiterpene lactones in rodents, J Ethnopharmacol, 128(1), 63-70.
27. Vandenbogaerde A *et al*., (2000), Evidence that total extract of Hypericum perforatum affects exploratory behaviour and exerts anxiolytic effects in rats, Pharmacol Biochem Behav, 65(4), 627-33.
28. Zanoli P, Avallone R, Baraldi M (2000), Behavioral characterisation of the flavonoids apigenin and chrysin, Fitoterapia, 71(1), S117-23.
29. Reginatto FH *et al*., (2006), Evaluation of anxiolytic activity of spray dried powders of two South Brazilian Passiflora species, Phytother Res, 20(5), 348-51.
30. Herrera-Ruiz M *et al*., (2008), Flavonoids from Tilia americana with anxiolytic activity in plus-maze test, J Ethnopharmacol, 118(2), 312-7.
31. Sampath C, Holbik M, Krenn L, Butterweck V (2011), Anxiolytic effects of fractions obtained from Passiflora incarnata L. in the elevated plus maze in mice, Phytother Res, 25(6), 789-95.
32. Roychoudhury M, Kulkarni SK (1997), Antianxiety profile of ondansetron, a selective 5-HT3 antagonist, in a novel animal model, Methods Find Exp Clin Pharmacol, 19(2), 107-11.
33. Raupp IM *et al*., (2008), Anxiolytic-like effect of chronic treatment with Erythrina velutina extract in the elevated plus-maze test, J Ethnopharmacol, 118(2), 295-9.
34. Wei XY, Yang JY, Wang JH, Wu CF (2007), Anxiolytic effect of saponins from Panax quinquefolium in mice, J Ethnopharmacol, 111(3), 613-8.
35. Viola H *et al*., (1994), Isolation of pharmacologically active benzodiazepine receptor ligands from Tilia tomentosa (Tiliaceae), J Ethnopharmacol, 44(1), 47-53.
36. Seo JJ *et al*., (2007), Anxiolytic-like effects of obovatol isolated from Magnolia obovata: involvement of GABA/benzodiazepine receptors complex, Prog Neuropsychopharmacol Biol Psychiatry, 31(7), 1363-9.
37. Rabbani M, Sajjadi SE, Jafarian A, Vaseghi G (2005), Anxiolytic effects of Salvia reuterana Boiss. on the elevated plus-maze model of anxiety in mice, J Ethnopharmacol, 101(1-3), 100-3

38. Awad R et al., (2003), Phytochemical and biological analysis of skullcap (Scutellaria lateriflora L. a medicinal plant with anxiolytic properties, Phytomedicine, 10(8), 640-9.
39. Guaraldo L, Chagas DA, Konno AC, Korn GP, Pfiffer T, Nasello AG (2000), Hydroalcoholic extract and fractions of Davilla rugosa Poiret: effects on spontaneous motor activity and elevated plus-maze behaviour, J Ethnopharmacol, 72(1-2), 61-7.
40. Mora S *et al*., (2005), Central nervous system activity of the hydroalcoholic extract of Casimiroa edulis in rats and mice, J Ethnopharmacol, 97(2), 191-7.
41. Kumar V, Jaiswal AK, Singh PN, Bhattacharya SK (2000), Anxiolytic activity of Indian Hypericum perforatum Linn: an experimental study, Indian J Exp Biol, 38(1), 36-41.
42. Cline M, Taylor JE, Flores J, Bracken S, McCall S, Ceremuga TE (2008), Investigation of the anxiolytic effects of linalool, a lavender extract, in the male Sprague-Dawley rat, AANA J, 76(1), 47-52.
43. Clénet F, Bouyon E, Hascoët M, Bourin M (2006), Light/dark cycle manipulation influences mice behaviour in the elevated plus maze, Behav Brain Res, 166(1), 140-9.
44. Boerngen-Lacerda R, Souza-Formigoni ML (2000), does the increase in locomotion induced by ethanol indicate its stimulant or anxiolytic properties, Pharmacol Biochem Behav, 67(2), 225-32.
45. Higgs S, Cooper SJ (1996), Hyperphagia induced by direct administration of midazolam into the parabrachial nucleus of the rat, Eur J Pharmacol, 313(1-2), 1-9.
46. Magee WJ, Eaton WW, Wittchen HU. Agoraphobia, simple phobia, and social phobia in the national comorbidity survey. Arch Gen Psychiatry, 1996; 53:159-68.
47. Kessler RC, Sonnega A, Bromet E. Post traumatic stress disorder in the national comorbidity survey. Arch Gen Psychiatry, 1995; 52:1048-60.
48. Rothbaum BO, Ninan PT, Thomas L. Sertraline in the treatment of rape victims with post traumatic stress disorder. J Trauma Stress, 1996; 9:865-71.
49. Schweitzer E. Generalized anxiety disorder: longitudinal course and pharmacologic treatment. Psychiatric Clin of N America, 1995; 18:843-57.
50. Roy-Byrne PP, Katon W. Generalized anxiety disorder in primary care: the precursor/modifier pathway to increased health care utilization. J Clin Psychiatry, 1997; 58 (Suppl 3): 34-8.

51. Markowitz JS, Weissman MM, Ouellette R, Lish JD, Klerman GL. Quality of life in panic disorder. Arch Gen Psychiatry, 1989;46:984-92.
52. Eddy MF, Walbroehl GS. Recognition and treatment of obsessive-compulsive disorder. Am Family Physician, 1998;57:1623-8, 1632-4.
53. Reecha Madaan, Anupam Sharma (2011), Evaluation of Anti-anxiety Activity of Actaea spicata Linn. International Journal of Pharmaceutical Sciences and Drug Research, 3(1), 45-47.
54. Agarwal MN, Itankar PR, Patil AT, Vyas JC, Kelkar AJ (2009), Anxiolytic activity of *Annona squamosa* leaf extracts in mice, Ind J Pharm Educ Res, 43(1), 99-103.
55. Kulkarni SK (1999), Handbook of experimental pharmacology, 3rd Ed. Vallabh Prakashan, New Delhi, 28-39.
56. Yadav AV, Kawale LA, Nade VS (2008), Effect of *Morus alba L.* (mulberry) leaves on anxiety in mice, Ind J Pharmacol, 40(1), 32-36.
57. Murali Madhav S (2001), Epidemiological study of prevalence of mental disorders in India, Ind J Comm Med, 26(4), 198-200.
58. Masoumeh E, Mohammad K, Maryam FA (2005), *Coriandrum sativum*: evaluation of its anxiolytic effects in the elevated plus maze, J Ethnopharmacol, 96, 365-370.
59. Eisenberg DM, David RB, Ettner SL (1998), Trends in alternative medicine use in the United states, J Am Med Assoc, 280, 1569-1575.
60. Parle M, Vasudevan M (2007), Memory enhancing activity of *Abana,* an Indian ayurvedic poly-herbal formulation, J Health Sci, 53(1), 43-52.
61. Plant extracts for treating skin disorders and enhancing healing of wounds for diabetic patients US8247004 B2 http://www.google.com/patents/US8247004
62. Janmejai K Srivastava, Eswar Shankar and Sanjay Gupta, Chamomile: A herbal medicine of the past with bright future, Mol Med Report. 2010; 3(6): 895–901.
63. Patel B. A., Patel P. U., Patel R. K. (2010), Physicochemical and phytochemical investigations of seeds of Celosia argentea Linn. International Journal of Pharmaceutical and Applied Sciences, Vol.1 (1), pp. 124-126.
64. OECD/OCDE (2008), OECD Guidelines For The Testing Of Chemicals, Acute Oral Toxicity – Up-and-Down-Procedure (UDP), 425, pp. 1-27.
65. Vogel H. G. (1996), Drug Discovery and Evaluation Pharmacological assay, Introduction-Strategies in drug discovery and evaluation, Springer-Verlag Berlin Heidelberg New York, 2nd edition, pp. 1-20.

66. Kulkarni SK. Handbook of experimental pharmacology. 3rd Ed. Vallabh Prakashan, New Delhi, 1999: 28-39.